

LPTENS-91/19

July, 1991

Classical W_n -Symmetry and Grassmannian Manifold

Y. Matsuo

Laboratoire de Physique Théorique

Ecole Normale Supérieure¹,

24 rue Lhomond, 75231 Paris CEDEX 05, France

Abstract

The classical W_n -symmetry is globally parametrized by the Grassmannian manifold which is described by the nonrelativistic fermions. We give the bosonization rule which defines the natural higher coordinates system to describe the W_n -geometry. The generators of the W_n -algebra can be obtained from the τ -function by using the vertex operator.

¹Unité propre du Centre National de la Recherche Scientifique, associée à l'Ecole Normale Supérieure et à l'Université de Paris-Sud

1 Introduction

Recently, a lot of attentions have been attracted to the investigation of the geometrical structure of the so-called W_n -gravity or the W_n -geometry.[1]-[8] This problem is challenging because it provides the natural extension of the two dimensional gravity and it will possibly be related to the higher dimensional field theories.[9] Although there are many works which discussed on such geometry, it is still far from the complete understanding because of its nonlinearity.

In this paper, we remark the relation between the global structure of the classical W_n -symmetry and the Grassmannian manifold. It is known[10] that the W_n -symmetry is closely related to the KdV equations.[11][12] Both of them are described as the deformation of rank n differential equations and the space of differential operators is parametrized by the Grassmannian manifold. In the KdV equations (or KP hierarchy), the nonlinear equations are mapped to linear motions on the Universal Grassmannian Manifold (UGM).[11] Our idea is that non-linear structure of W_n symmetry will be simplified also in the language of the Grassmannian. Although we have not fully completed this program, the mere introduction of the Grassmannian seems to provide many clues to understand the geometry. It is known that the natural tool to describe the Grassmannian manifold is the free fermion theory. Unlike the KP hierarchy theory, our fermions turns out to be non-relativistic. It reflects the fact that the particle number is always finite. We show that the deformation of n -th order differential operators can be linearly described by the bosonization. We prove that the bosonic degree of freedom is finite dimensional (say $t_1, \dots, t_n$) and the first one (t_1) corresponds to the holomorphic coordinate of the Riemann surface.

The most important merit to introduce such higher coordinates is that W_n -algebra generators can then be obtained from a single generator, i.e. the τ -function. In this sense, the situation becomes similar to that of the super Virasoro symmetry with the supercoordinate.

2 Classical W_n -symmetry

In various contexts, the classical W_n -symmetry is realized by the deformation of n -th order differential equation,[1][2][7][8]

$$\mathcal{L}_n f(t) = 0, \quad (1)$$

$$\mathcal{L}_n = \partial^n + u_1(t)\partial^{n-1} + u_2(t)\partial^{n-2} + \cdots u_n(t). \quad (2)$$

We call the coefficient functions $u_s(t)$ gauge potentials in the following. In order to get the W_n symmetry, we need to require the constraint for the first gauge potential,

$$u_1(t) \equiv 0. \quad (3)$$

The deformation is generated by a pair of differential operators X and Y ². We need to choose them such that $\mathcal{L}_n$ keeps its form (2,3) under the variation of the following type,

$$\delta_X \mathcal{L}_n \equiv Y \mathcal{L}_n - \mathcal{L}_n X \quad (4)$$

$$= \delta_X u_2 \partial^{n-2} + \cdots + \delta_X u_n, \quad (5)$$

$$\delta_X f = X f(t) \quad (6)$$

The relation between the operators X, Y is important. From eq.(4), Y is formally given by

$$\begin{aligned} Y &= \mathcal{L}_n X \mathcal{L}_n^{-1} + \delta \mathcal{L}_n \mathcal{L}_n^{-1} \\ &= (\mathcal{L}_n X \mathcal{L}_n^{-1})^+. \end{aligned} \quad (7)$$

Here we used the projection of the pseudo differential operator into positive parts $(\)^+$. Explicitly, if $O = \sum_{\ell=0}^{\infty} o_\ell \partial^{n-\ell}$, we get $(O)^+ = \sum_{\ell=0}^n o_\ell \partial^{n-\ell}$. We define also $(O)^- = O - (O)^+$. If we ignore the constraint (3), we can start from any differential operator X to get the deformation of $\mathcal{L}_n$,

$$\delta_X \mathcal{L}_n = \mathcal{B}_X \mathcal{L}_n, \quad \mathcal{B}_X = -(\mathcal{L}_n X \mathcal{L}_n^{-1})^-. \quad (8)$$

We remark that although the deformation equation (8) reminds us of that of the KP hierarchy,[11] there are some important differences. In the KP hierarchy, we define the 0-th order pseudo-differential operator (PSD) $S = 1 + s_1(t)\partial^{-1} + s_2(t)\partial^{-2} + \cdots$ through $\mathcal{L}_n = S\partial^n S^{-1}$. Such operator can

² We use the notation of the ref. [8]

be determined up to right multiplication of constant 0-th order PSD C , $S \rightarrow SC$. Then “time evolution” of the system is governed by $\frac{\partial}{\partial t_\ell} S = A_\ell S$ with $A_\ell = -(S\partial^\ell S^{-1})^-$. These time evolutions are then interpreted as linear motions on the Universal Grassmannian Manifold which is directly related to S . This formulation has a merit that we can discuss the deformation of the differential operators of any order in the same framework. However, the relation between s_ℓ and gauge potentials is nonlinear. Moreover, the Poisson bracket structure of $\mathcal{L}_n$ is written in terms of the variation (8). If we the inner product $(A, L_n) = \int dt \sum_{\ell=1}^n a_\ell(t) u_\ell(t)$ for $A = \sum_{\ell=1}^n \partial^{\ell-1-n} a_\ell(t)$, the second Poisson bracket[12] is written by,

$$\{(A, \mathcal{L}_n), (B, \mathcal{L}_n)\} = -(B, (\mathcal{L}_n A)^- \mathcal{L}_n). \quad (9)$$

The RHS of this equation is the variation of $\mathcal{L}_n$ with $X = (A\mathcal{L}_n)^+$. This gives the classical definition of the W_n algebra. Hence in order to understand its geometrical structure, we had better introduce the Grassmanian which is directly related to $\mathcal{L}_n$ itself.

3 Grassmannian manifold

The correspondence between n -th order differential equation and the Grassmannian is more or less known through the essential role of the Wronskian. In our paper, we make it in more explicit form and develop the suitable machinery to describe it.

Define a Hilbert space $\mathcal{H}$ as the space of functions which is regular at $t = 0$, i.e. $f \in \mathcal{H}$ if it has the Taylor expansion,

$$f(t) = \sum_{n=0}^{\infty} \frac{f_n}{n!} t^n. \quad (10)$$

Suppose all the coefficients in $\mathcal{L}_n$ are regular at $t = 0$ ³. Then the kernel the differential operator (1) determines an n -dimensional subspace of $\mathcal{H}$. Denote the Grassmannian manifold which consists of n -dimensional subspaces of $\mathcal{H}$ by $Gr^{(n)}$. The correspondence discussed above shows that there is a mapping

³ If we allow that the gauge potentials might have singularities at the origin $t = 0$, we need to extend the Hilbert space to $\mathbf{C}[[\log t]]$ instead of $\mathbf{C}[[t]]$. We must modify our discussion in the following if we need to discuss this generalized situation.

from the space of n -th order differential operators into the Grassmanian $Gr^{(n)}$. It is important to notice that there is the inverse mapping. Let Ω be a point of $Gr^{(n)}$. Pick up a basis of Ω , $\{f_1, \dots, f_n\}$. We can get n -th order differential equation,

$$\mathcal{L}_n(\Omega)f = \tau_n(\Omega)^{-1} \begin{vmatrix} f & f_1 & \cdots & f_n \\ \partial f & \partial f_1 & \cdots & \partial f_n \\ \vdots & \vdots & & \vdots \\ \partial^n f & \partial^n f_1 & \cdots & \partial^n f_n \end{vmatrix} \quad (11)$$

where, $\mathcal{L}_n(\Omega) \equiv \partial^n + u_1(\Omega)\partial^{n-1} + \dots + u_n(\Omega)$ and

$$\tau_n(\Omega) \equiv \begin{vmatrix} f_1 & \cdots & f_n \\ \vdots & & \vdots \\ \partial^{n-1} f_1 & \cdots & \partial^{n-1} f_n \end{vmatrix}. \quad (12)$$

In this way, one observe the bijection between the Grassmanian $Gr^{(n)}$ and the space of the n -th order differential operators (without u_1 constraint (3)). In a sense, the functions $\{u_1(\Omega), \dots, u_n(\Omega)\}$, can be regarded as inhomogeneous coordinates of $Gr^{(n)}$.

Now let us come back to the classical W_n -symmetry. It is defined as the deformations of $\mathcal{L}_n$. Since the Grassmannian $Gr^{(n)}$ parametrizes the family of differential operators, W_n -algebra generators can be regarded as the elements of the tangent space of the Grassmannian, $T_*Gr^{(n)}$ ⁴. In this sense, the Grassmannian gives the global structure of the W_n -algebra. The merit to lay emphasis on the Grassmannian is that it is a homogenous space and basically a linear object. As we are going to discuss, we can treat it neatly through the free fermion operator formalism and its bosonization.

4 Free fermion formalism

As in the KP hierarchy, the free fermion operator formalism provides a systematic and powerful tool to analyze the structure of the Grassmannian. In order to apply to our case, however, we need to make a slight modification.

⁴ Precisely, we need to impose the constraint to get the W_n symmetry. It will be discussed in the last section.

It is due to the fact that the Wronskians which appeared in the previous section had finite ranks. Correspondingly, we need to deal with the states with finite particle number. This situation is similar to the free fermion theory of the finite N matrix models.[13] In fact, the nonrelativistic fermion theory used there can be also applied to our case.

Let us define the free fermion operators and the Fock vacuum as in the matrix model.

$$\{\psi_n, \psi_m\} = \{\psi_n^*, \psi_m^*\} = 0, \quad (13)$$

$$\{\psi_n, \psi_m^*\} = \delta_{n,m}, \quad (n, m = 0, 1, \dots) \quad (14)$$

$$\psi_n^* |\emptyset\rangle = 0 \quad \forall n, \quad (15)$$

$$\langle \emptyset | \psi_n = 0 \quad \forall n. \quad (16)$$

Note that we use the semi-infinite indices $n = 0, 1, 2, \dots$ for the fermion operators. The vacuum states $|\emptyset\rangle$ and $\langle \emptyset|$ correspond to the no particle states. The n -particle ground state can be created from them in the standard way,

$$|n\rangle = \psi_{n-1} \psi_{n-2} \cdots \psi_0 |\emptyset\rangle, \quad (17)$$

$$\langle n| = \langle \emptyset | \psi_0^* \psi_1^* \cdots \psi_{n-1}^*. \quad (18)$$

The $U(1)$ current operators,

$$J_n = \sum_{s=0}^{\infty} \psi_{n+s} \psi_s^*, \quad (19)$$

shall play the role of the ‘‘Hamiltonian’’ as it does in the KP hierarchy.

We represent the Hilbert space $\mathcal{H}$ by the ‘‘one-particle state’’ in the fermion Fock space. For each $f(t) = \sum_n \frac{f_n}{n!} t^n \in \mathcal{H}$, define $\psi_f^* \equiv \sum_n f_n \psi_n^*$. We can show,

$$f(t) = \langle \emptyset | \psi_f^* \exp(J_1 t) | 1 \rangle. \quad (20)$$

More generally, we can represent $Gr^{(n)}$ in our Fock space. For each $\Omega \in Gr^{(n)}$, choose a frame $\{f_1, \dots, f_n\}$. It is mapped into the n -particle states $\langle \Omega| \equiv \langle \emptyset | \psi_{f_1}^* \cdots \psi_{f_n}^*$. This procedure has an ambiguity up to the multiplication of the constant number which depends on the choice of the frame. This is the standard Plücker embedding⁵. It is easily proved that the determinant $\tau_n(\Omega)$

⁵ Conversely, generic points of the n -particle states do not correspond to the Grassmannian. The Grassmannian points are characterized by the Hirota’s bilinear identity, $\sum_{s=0}^{\infty} \langle \Omega | \psi_s \otimes \langle \Omega | \psi_s^* = 0$.

(12) can be represented as the vacuum expectation value (VEV) of n -particle states,

$$\tau_n(\Omega) = \langle \Omega | \exp(J_1 t) | n \rangle. \quad (21)$$

The differential equation is now written simply as $\langle \Omega | \psi_f^* = 0$. Instead of using this form, it is more convenient to introduce following “generating functional” for the gauge potentials,

$$\begin{aligned} W_\Omega(\zeta) &= \sum_{s=0}^n \zeta^s u_s(\Omega) \tau_n \\ &= \langle \Omega | \exp(J_1 t) \psi^*(\zeta) | n+1 \rangle. \end{aligned} \quad (22)$$

with $\psi^*(\zeta) = \sum_{n=0}^{\infty} \psi_n^* \zeta^n$. In order to prove these identities, we used a lemma, $\partial^n f(t) = \langle \emptyset | \psi_f \exp(t J_1) \psi_n | \emptyset \rangle$. In the terminology of the KP hierarchy, τ_n and $W_\Omega(\zeta)$ correspond to the τ -function and the Baker-Akhiezer function respectively. More generally, we define the generating functional of the character functions,

$$W_\Omega(\zeta_1, \dots, \zeta_m) = \langle \Omega | \exp(J_1 t) \psi^*(\zeta_1) \cdots \psi^*(\zeta_m) | n+m \rangle. \quad (23)$$

5 Bosonization

We would like to discuss how the deformation of the differential equation looks like in our fermion formalism. The natural transformation which acts on the Grassmannian is the general linear transformation of $\mathcal{H}$, i.e. so called $w_{1+\infty}$ algebra.[15] In the free fermion language, the transformation is generated by the fermion bilinear,

$$w_{n,m} = \sum_{s=m}^{\infty} \frac{s!}{(s-m)!} \psi_{s+n-m} \psi_s^*. \quad (24)$$

The commutation relation with $\psi^*(\zeta)$ illuminates the origin of this operator,

$$[w_{n,m}, \psi^*(\zeta)] = -\zeta^n \frac{\partial^m}{\partial \zeta^m} \psi^*(\zeta). \quad (25)$$

When it applied to the one particle state it induces the $w_{1+\infty}$ transformation,

$$\langle \emptyset | \psi_f^* w_{n,m} \exp(t J_1) | 1 \rangle = t^m \frac{\partial^n}{\partial t^n} f(t). \quad (26)$$

We should remark that the indices n and m are interchanged. It is because the Hamiltonian $\exp(tJ_1)$ actually plays the role of Laplace transformation here. Similar phenomena also happened in the finite N matrix models.

Now the variation of the n -particle bra state $\langle \Omega|$ is simply given by, $\delta_{n,m} \langle \Omega| = \langle \Omega| w_{n,m}$. We need to rewrite its action in terms of the bosonic variable. However, in order to describe the multi-particle states, the single holomorphic coordinate t is not enough. As in the KP hierarchy, we need to introduce the infinite number of time coordinates t_n ($n = 0, 1, 2, \dots$) through the modification of the Hamiltonian,

$$\exp(tJ_1) \rightarrow \exp\left(\sum_{\ell=0}^{\infty} t_{\ell} J_{\ell}\right) \equiv \exp(H([t])). \quad (27)$$

Although we shall show later that only n coordinates are enough to describe n -particle states, it is convenient to keep all of them for a while.

The bosonization of the non-relativistic fermion in the context of the matrix model was studied previously by Itoyama and current author.[14] We can copy it to apply to this case. Define $w_n(\zeta) \equiv \sum_{m=0}^{\infty} w_{m,n} \zeta^{-m-1}$. The key identity⁶ is,

$$w_n(\zeta) |k\rangle = \frac{1}{n+1} Q_{n+1}(w_0(\zeta)) |k\rangle. \quad (28)$$

with

$$Q_n(f) = (\partial_{\zeta} + f(\zeta))^n \cdot 1 = e^{-F(\zeta)} \partial^n e^{F(\zeta)}, \quad (\partial F \equiv f). \quad (29)$$

We shall prove later that $Q_{\ell}(w_0(\zeta)) |k\rangle = 0$ identically if $\ell > k$. We also use the following lemma,

$$w_n(\zeta) \exp(H[t]) = \exp(H([t])) \left(\sum_{\ell=0}^n {}_n C_{\ell} Q_{n-\ell}(j(\zeta)) w_{\ell}(\zeta) \right)_{-}, \quad (30)$$

where $(\)_{-}$ is the projection to the negative power of ζ . In the following, we use the notations, $j(\zeta) = \sum_{\ell=1}^{\infty} \ell t_{\ell} \zeta^{\ell-1}$. and $\frac{d}{dj} = \sum_{\ell=0}^{\infty} \zeta^{-\ell-1} \frac{\partial}{\partial t_{\ell}}$.

⁶ We need a “translation” from identities obtained in [14]. It is important to notice that,

$$\begin{aligned} \det(\lambda_i^j)_{i,j=1,\dots,n} &= \langle \emptyset | \psi^*(\lambda_1) \cdots \psi^*(\lambda_n) | n \rangle, \\ \langle \emptyset | \psi^*(\lambda_1) \cdots \psi^*(\lambda_n) w_{p,q} &= \sum_{i=1}^n \lambda_i^p \frac{\partial^q}{\partial \lambda_i^q} \langle \emptyset | \psi^*(\lambda_1) \cdots \psi^*(\lambda_n). \end{aligned}$$

Combining these identities, we get the transformations of the τ -function in the bosonized form,

$$\delta_s(\zeta)\tau_n = \tilde{\delta}_s(\zeta)\tau_n, \quad (31)$$

with

$$\tilde{\delta}_s(\zeta) = \sum_{\ell=1}^{s+1} C_\ell \left(Q_{s+1-\ell}(j(\zeta)) Q_\ell \left(\frac{d}{dj}(\zeta) \right) \right)_-. \quad (32)$$

Here, we use the notation $(\)_-$ which means the projection to the negative powers of ζ .

Eq.(31) gives us very important corollary. Since every $w_{1+\infty}$ generators are bosonized, we can get more direct “bosonization” rules. The most useful one is that

$$\exp(H([t]))\psi^*(\zeta)|n+1> = V_n^*(\zeta)\exp(H([t]))|n>, \quad (33)$$

with,

$$V_n^*(\zeta) = \zeta^n \exp \left(- \sum_{\ell=1}^{\infty} \frac{\zeta^{-\ell}}{\ell} \frac{\partial}{\partial t_\ell} \right). \quad (34)$$

This relation explicitly shows that we can get $W_\Omega(\zeta)$ out of τ -function, $W_\Omega(\zeta) = V_n^*(\zeta)\tau_n([t])$. It seems rather nontrivial that the RHS is a n -th polynomial in terms of ζ . We shall explicitly prove it later. Anyway, the importance of this formula is that it implies that the W_n charges can be obtained from τ -function in a simple way. In this sense, we can derive all the properties of W_n algebra in terms of the single function τ_n .

Our method is so systematic that we can calculate the transformation rules of the character polynomials without any difficulty. We write some of the useful identities,

$$\begin{aligned} [w_n(\zeta_1), \psi^*(\zeta_2)] &= -\frac{1}{\zeta_1 - \zeta_2} \frac{\partial^n}{\partial \zeta_2^n} \psi^*(\zeta_2), \\ \psi^*(\zeta_1) \cdots \psi^*(\zeta_n) Q_\ell(w_0(\zeta)) &= Q_\ell(w_0 + \sum_{s=1}^m \frac{1}{\zeta - \zeta_s}) \psi^*(\zeta_1) \cdots \psi^*(\zeta_m) \\ Q_\ell(\sum_{s=1}^m \frac{1}{\zeta - \zeta_s}) &= \prod_{s=1}^m (\zeta - \zeta_s)^{-1} \partial_\zeta^\ell \prod_{s=1}^m (\zeta - \zeta_s), \quad \text{etc...} \end{aligned}$$

For example, the transformation rule of $W_\Omega(\zeta)$ reads,

$$\delta_s(\zeta_1)W_\Omega(\zeta_2) = \tilde{\delta}_s(\zeta_1, \zeta_2)W_\Omega(\zeta_2), \quad (35)$$

with,

$$\begin{aligned}\tilde{\delta}_s(\zeta_1, \zeta_2) &= \sum_{\ell=0}^s {}_s C_{\ell} \left(\frac{1}{\ell+1} Q_{s-\ell}(j(\zeta_1)) Q_{\ell+1}\left(\frac{d}{dj}(\zeta_1)\right) \right)_{-} \\ &\quad + \frac{{}_s C_{\ell}}{\zeta_1 - \zeta_2} \left(Q_{s-\ell}(j(\zeta_1)) Q_{\ell}\left(\frac{d}{dj}(\zeta_1)\right) - Q_{s-\ell}\left(\frac{d}{dj}(\zeta_1)\right) \frac{\partial^{\ell}}{\partial \zeta_2^{\ell}} \right)_{-}\end{aligned}\quad (36)$$

Generalizations to the higher polynomials are straightforward.

6 Effective time coordinates

As we have remarked before, only finite number of “times” are needed to describe the Grassmanian $Gr^{(n)}$. The simplest situation is the one-particle case, where we can simply replace $\frac{\partial}{\partial t_s}$ by $\frac{\partial^s}{\partial t_1^s}$. For n -particle states, the situation is more complicated. Note that the n -particle state can be represented as the n -th wedge product of the single particle wave functions, $\Omega = f_1 \Lambda \cdots \Lambda f_n$. Let us denote the differentiation on the α -th wave function as $\frac{\partial}{\partial t_s}(\alpha)$. Then it is obvious that

$$\frac{\partial}{\partial t_s} = \sum_{\alpha=1}^n \frac{\partial}{\partial t_s}(\alpha) = \sum_{\alpha=1}^n \left(\frac{\partial}{\partial t_1}(\alpha) \right)^s \quad (37)$$

The problem reduces to finding algebraically independent basis for the symmetric function of n -variables. The desired basis is clearly $\Gamma_n = \left\{ \frac{\partial}{\partial t_{\ell}} \right\}_{\ell=1,2,\dots,n}$. The explicit expression of $\frac{\partial}{\partial t_{s+n}}$ in terms of Γ is more involved. However, it is fortunate that these derivatives appear only through the combinations, $Q_{\ell}\left(\frac{d}{dj}(\zeta)\right) = (V_n^*(\zeta))^{-1} \partial_{\zeta}^{\ell} V_n^*(\zeta)$. Hence we need to express only V_n^* in terms of the basis Γ_n . On n -particle states, t_0 derivative is simply replaced by n . ($J_0|n\text{-particle}\rangle = n|n\text{-particle}\rangle$.) Hence,

$$\begin{aligned}V_n^* &= \zeta^n \exp \left(- \sum_{\ell=1}^{\infty} \frac{1}{\ell} \zeta^{-\ell} \frac{\partial}{\partial t_{\ell}} \right) \quad (n\text{-particle}) \\ &= \zeta^n \prod_{\alpha=1}^n \exp \left(- \sum_{\ell=1}^{\infty} \frac{1}{\ell} \zeta^{-\ell} \frac{\partial^{\ell}}{\partial t_1^{\ell}}(\alpha) \right) \\ &= \zeta^n \prod_{\alpha=1}^n \left(1 - \frac{1}{\zeta} \frac{\partial}{\partial t_1}(\alpha) \right)\end{aligned}\quad (38)$$

Here, there appear only up to n -th order symmetric polynomial in V_n^* . They can be explicitly expressed in terms of Γ_n . Let us introduce polynomials $p_n([c])$ as follows,

$$\sum_{\ell=0}^{\infty} p_{\ell}([c]) \zeta^{\ell} = \exp \left(\sum_{\ell=1}^{\infty} c_{\ell} \zeta^{\ell} \right). \quad (39)$$

For example, $p_0 = 1$, $p_1 = c_1$, $p_2 = \frac{1}{2}c_1^2 + c_2$, $\dots$. p_n is written out of $c_1, \dots, c_n$. We claim that

$$V_n^*(\zeta) = \sum_{\ell=0}^n \zeta^{n-\ell} p_{\ell}([c])|_{c_s = -\frac{1}{s} \frac{\partial}{\partial t_s}}. \quad (40)$$

Note that V_n^* is an n -th order polynomial in terms of ζ . This proves explicitly that $Q_{\ell}(w_0(\zeta))|n\text{-particle} \rangle = 0$, if $\ell > n$. This fact was mentioned before.

7 Discussions

To summarize, the deformation of the n -th order differential equation is understood as the analogue of the Bäcklund transformation in the KP hierarchy. Its transformation is written by using a kind of boson-fermion correspondence. The main difference between the two theory is that the number of time coordinates is finite in our case. The natural question is what is the analogue of the KP equation itself. The answer is straightforward. Let us think the deformation induced by the derivative of a higher coordinate t_s , $\delta f = \frac{\partial}{\partial t_s} f$ in (1). Since $\frac{\partial}{\partial t_s} = \left(\frac{\partial}{\partial t_1} \right)^s$ on the one-particle state, we can take $X = \left(\frac{\partial}{\partial t_1} \right)^s$. The deformation of $\mathcal{L}_n$ follows directly from (8),

$$\frac{\partial}{\partial t_s} \mathcal{L}_n = \mathcal{B}_s \mathcal{L}_n, \quad \mathcal{B}_s \equiv -(\mathcal{L}_n \partial_1^s \mathcal{L}_n^{-1})^-. \quad (41)$$

If we start from the UGM defined through S , we would get the higher KdV equations,

$$\frac{\partial}{\partial t_s} \mathcal{L}_n = [A_s, \mathcal{L}_n], \quad A_s = (\mathcal{L}_n^{s/n})^+. \quad (42)$$

We should not confuse these two nonlinear equations.

Throughout this letter, we do not pay enough attention to the $U(1)$ -constraint (3). In our framework, this constraint does not arise naturally

and we need to impose it by hand. In terms of τ -function, this constraint is given by

$$\partial_1 \tau([t])|_{t_2=t_3=\dots=0} = 0. \quad (43)$$

Although it is easy to obtain known formulae in our language, we have not clarified the geometrical origin of this constraint.

In the subsequent paper, we shall discuss the Hamiltonian structure in terms of τ -function and discuss the free field realization. It will be also interesting to discuss the $N = 2$ supersymmetric extension of W_n symmetry[16] because it is formulated almost parallel to the bosonic case.

Acknowledgements: The author would like to thank J.L. Gervais for the discussions about the classical W_n -symmetry and the Toda field theory. He is also obliged to V.Fateev for his important comments and encouragements.

References

- [1] A.B.Zamolodchikov, *Theor. Math. Phys.* **65** (1985) 1205; V.A. Fateev and A.B.Zamolodchikov, *Nucl. Phys.* **B280** [FS18] (1987) 644; V.A. Fateev and S.L. Lukyanov, *Int. J. Mod. Phys.* **A3** (1988) 507; V.A. Fateev and S.L. Lukyanov, "Poisson-Lie Groups and Classical W-Algebra", preprint Paris Par-LPTHE 91-16, (February, 1991).
- [2] A. Bilal and J.-L. Gervais, *Phys. Lett.* **B206** (1988) 412; *Nucl. Phys.* **B314** (1989) 646; *Nucl. Phys.* **B318** (1989) 579.
- [3] Y. Matsuo *Phys. Lett.* **227B** (1989) 222; in proceeding of the meeting *Geometry and Physics*, Lake Tahoe, July 1989.
- [4] M. Bershadsky and H. Ooguri, *Comm. Math. Phys.* **126** (1989) 49.
- [5] K. Schoutens, A. Sevrin and P. van Nieuwenhuizen, *Phys. Lett.* **243B** (1990) 245; *Nucl. Phys.* **B349** (1991) 791; "Quantum W_3 Gravity in the Chiral Gauge" preprint ITP-SB-91-07, (March 1991).
- [6] C.M. Hull, *Phys.Lett.* **240B** (1990) 110.

- [7] G. Sotokov and M. Stanishkov, *Nucl. Phys.* **B356** (1991)439; G. Sotokov, M. Stanishkov and C. J. Zhu, *Nucl. Phys***B356** (1991) 245.
- [8] P.Di Francesco, C. Itzykson and J.B. Zuber, “Classical W-Algebras” preprint SPhT/90-149, PUPT-1211.
- [9] Q. H. Park, *Phys. Lett.* **B236** (1990) 429; K. Yamagishi and G. F. Chapline, *Class. Quantum Grav.* **8** (1991) 185; N.J. Hitchin *Prog. Theor. Phys. Suppl.* **102** (1990) 159.
- [10] J.-L. Gervais, *Phys. Lett.* **B160** (1985) 277; K. Yamagishi, *Phys. Lett* **B205** (1988) 406; I. Bakas, *Phys. Lett.* **B213** (1988) 313.
- [11] M. Sato, *RIMS Kokyuroku* **439** (1981) 30; E. Date, M. Jimbo, M.Kashiwara and T.Miwa, “Transformation Groups for Soliton Equations” in *Proc. of RIMS Symposium on Non-lineq Integrable Systems-Classical Theory and Quantum Theory* (Kyoto, Japan, May 1981) (World Scientific Publication Co. Singapore, 1983); G.Segal and G.Wilson, *Pub. Math. IHES* **61** (1985) 5.
- [12] V.G. Drinfeld and V.V. Sokolov, *Journ. Sov. Math.* **30** (1985) 1975.
- [13] T. Banks, M.Douglas, N.Seiberg and S.Shenker, *Phys. Lett.* **B238** (1990) 279.
- [14] H. Itoyama and Y. Matsuo, *Phys.Lett.* **B255** (1991) 202; *Phys. Lett.* **B262** (1991) 233.
- [15] I. Bakas, *Phys. Lett.* **B228** (1989) 57; C. Pope, I. Romas and X. Shen, *Nucl. Phys.* **B339** (1990) 191; *Phys. Lett.* **B242** (1990) 401; *Phys. Lett.* **B245** (1990) 72; M. Fukuma, H.Kawai and R. Nakayama, “Infinite Dimensional Grassmanian Structure of Two-Dimensional Quantum Gravity” Tokyo preprint (Nov. 1991).
- [16] T.Inami and H. Kanno, “ $N = 2$ Super W algebra and Generalized $N = 2$ Super KdV Hierarchies Based on Lie Superalgebras” preprint YITP/K-928 (May 1991).